Total number of printed pages-11

3 (Sem-2/CBCS) CHE HC 2



2022

CHEMISTRY

(Honours)

Paper: CHE-HC-2026

(Physical Chemistry-II)

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

- 1. Answer **any seven** of the following questions: $1 \times 7 = 7$
 - (a) Give the SI unit of energy.
 - (b) Define specific heat of a system.

- The variation of enthalpy of a reaction with temperature is given by
 - Hess's law
 - Kirchhoff's equation,
 - Henry's law, (iii)
 - Raoult's law

(Choose the correct option)

- A process is carried out at constant pressure and temperature. It will be spontaneous if
 - $\Delta G < 0$
 - (ii) $\Delta H < 0$
 - (iii) $\Delta U < 0$
 - $\Delta S < 0$ (iv)

(Choose the correct option)

- A solution is a
 - homogeneous mixture of only two components

- homogeneous mixture of any (ii) number of components
- heterogeneous mixture
- (iv) anything mixed with water (Choose the correct option)
- What is excess thermodynamic function?
- Name a colligative property that is used to determine the molar mass of a protein.
- Equimolar solutions of glucose and sodium chloride are not isotonic. Justify.
- Find the value of work done when 2 moles of an ideal gas is allowed to expand from 1 L to 10 L against vacuum at 298K.
- Name the thermodynamic property that measures the disorderliness of a system.

3 (Sem-2/CBCS) CHE HC 2/G 3 Contd.

- 2. Answer *any four* of the following questions: 2×4=8
 - (a) Define intensive property. Give one example.
 - (b) State Zeroth law of thermodynamics.
 - (c) Define explosion temperature and adiabatic maximum flame temperature.
 - (d) What do you mean by network? Briefly explain.
 - (e) Explain residual entropy.
 - (f) Define fugacity function.
 - (g) An ideal gas undergoes a single step expansion a constant external pressure P from (P_1, T, V_1) to (P, T, V_2) . What is the magnitude of work done by the system?

(h) Find ΔH of the reaction : $H_2(g) + Br_2(g) \longrightarrow 2HBr(g)$ Given : $\Delta H_{H-H} = 435.1, \Delta H_{Br-Br} = 192.5,$

 $\Delta H_{H-Br} = 368.2 \ kJ/mol.$

- 3. Answer *any three* of the following questions: 5×3=15
 - (a) (i) State Path function with suitable example.
 - (ii) Show that in an isothermal expansion, the work is done at the expense of the heat absorbed. 3
 - (b) Derive the Gibbs Helmholtz equation.
 - (c) (i) Write short note on the third law of thermodynamics.
 - (ii) Explain briefly how absolute entropy of a molecule can be determined from heat capacity measurement.

- (d) Give the criteria of spontaneity and thermodynamic equilibrium in terms of enthalpy, entropy, Helmholtz free energy and Gibbs free energy.
- (e) (i) Calculate $K_{\rm C}$ for the reaction $2SO_3(g) \Longrightarrow 2SO_2(g) + O_2(g) \quad \text{for}$ which $K_p = 3.5 \times 10^{-23}$ atm at 27° C.
 - (ii) How molar mass can be determined from freezing point depression?
- (f) (i) 0.5g of a non-volatile solute of molar mass 60g mol⁻¹ is dissolved in 100g of ethyl acetate at 20°C. What would be the vapour pressure of this solution at 20°C? The vapour pressure of ethyl acetate at 20°C is 72.8 Torr.
 - (ii) Explain briefly any one method for measurement of vapour pressure lowering.

- (g) What is osmotic pressure? Give detailed thermodynamic derivation of osmotic pressure of a solution having non-volatile solute.
- (h) What are colligative properties?

 Explain two practical applications of colligative properties.
- 4. Answer *any three* of the following questions: 10×3=30
 - (a) (i) State and explain first law of thermodynamics. Show that for isochoric process, $q = \Delta U$. 3+2=5
 - (ii) Derive the integrated Kirchhoff equation.
 - (b) (i) Define heat capacity of a system. Show that $C_p C_v = R$ for 1 mole of an ideal gas. 1+3=4
 - (ii) State and explain Raoult's law for vapour pressure of binary solution of volatile liquid. What is an ideal solution?

- (c) (i) Calculate q, w, ΔU and ΔH for the reversible isothermal expansion of one mole of an ideal gas at 27°C from a volume of 10 dm^3 to a volume of $20 dm^3$.
 - (ii) Explain that the entropy of the universe is increasing continuously.
 - (iii) Explain briefly the vapour pressure vs. composition diagram of a binary liquid mixtures having positive deviation.
- (d) (i) Explain that the thermodynamic isothermal reversible work of expansion is the maximum work.

(ii) Give the thermodynamic derivation of the relation between Gibb's free energy of a reaction and its reaction quotient.

3

- (iii) Give two limitations of first law of thermodynamics. 2
- (e) (i) Define enthalpy of neutralization.
 - (ii) The enthalpy of combustion of glucose $C_6H_{12}O_6(S)$ is -2816 $kJ \, mol^{-1}$ at 25°C. Calculate ΔH_f^o of $C_6H_{12}O_6(S)$. The ΔH_f^o values for $CO_2(g)$ and $H_2O(l)$ are -393.5 and -286.2 $kJ \, mol^{-1}$ respectively.
 - (iii) Give a brief account of coupling of exoergic and endoergic reactions.
 - (iv) State and explain van't Hoff theory of dilute solution as applied to osmotic pressure.
- (f) (i) Discuss about the molecular and statistical interpretation of entropy. $2\frac{1}{2} \times 2=5$

3 (Sem-2/CBCS) CHE HC 2/G 9

Contd.

- (ii) Show that : $\Delta G_{mix} = nRT \left(x_1 \ln x_1 + x_2 \ln x_2 \right) \qquad 5$
- (g) (i) Prove that : $\left(\frac{\partial V}{\partial T}\right)_P = -\left(\frac{\partial S}{\partial P}\right)_T$
 - (ii) Explain the variation of chemical potential with temperature. 3
 - (iii) Calculate the pressure of CO_2 gas at 700K in the heterogeneous equilibrium reaction $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ if ΔG^o for this reaction is $130.2 \ kJmol^{-1}$.
- (h) (i) Show that : $K_p = K_x (P)^{\Delta ng} = K_c (RT)^{\Delta ng}$ under what conditions, $K_p = K_x = K_c?$ 5+1=6

(ii) State and explain Le Chatelier's principle taking any one example.

4

