

Total number of printed pages-11

3 (Sem-3 /CBCS) MAT HC 1

2022

MATHEMATICS

(Honours)

Paper : MAT-HC-3016

(Theory of Real Functions)

Full Marks : 80

Time : Three hours

The figures in the margin indicate full marks for the questions.

1. Answer **any ten** parts : $1 \times 10 = 10$

(a) Is every point in I a limit point of $I \cap Q$?

(b) Find $\lim_{x \rightarrow 1} \frac{x^2 - x + 1}{x + 1}$.

(c) Let $f(x) = \text{sgn}(x)$. Write the limits

$\lim_{x \rightarrow 0^+} f(x)$ and $\lim_{x \rightarrow 0^-} f(x)$.

Contd.



(d) Let $p: \mathbb{R} \rightarrow \mathbb{R}$ be the polynomial function

$$p(x) := a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

if $a_n > 0$, then $\lim_{x \rightarrow \infty} p(x) = ?$

(e) Let f be defined on $(0, \infty)$ to \mathbb{R} . Then the statement

" $\lim_{x \rightarrow \infty} f(x) = L$ if and only if

$\lim_{x \rightarrow 0^+} f\left(\frac{1}{x}\right) = L$ " is true **or** false.

(f) Let $A \subseteq \mathbb{R}$ and let f_1, f_2, \dots, f_n be function on A to \mathbb{R} , and let c be a cluster point of A . If $\lim_{x \rightarrow c} f_k(x) = L_k$,

$k = 1, 2, \dots, n$,

then $\lim_{x \rightarrow c} (f_1 \cdot f_2 \cdot \dots \cdot f_n) = ?$

(g) Is the function $f(x) = \frac{1}{x}$ continuous on $A = \{x \in \mathbb{R} : x > 0\}$?

(h) Write the points of continuity of the function $f(x) = |x|$.

(i) "A rational function is continuous at every real number for which it is defined." Is it true or false?

(j) "Let f, g be defined on \mathbb{R} and let $c \in \mathbb{R}$. If $\lim_{x \rightarrow c} f(x) = b$ and g is continuous at b , then $\lim_{x \rightarrow c} (g \cdot f)(x) = g(b)$." Write whether this statement is correct or not.

(k) The functions $f(x) = x$ and $g(x) = \sin x$ are uniformly continuous on \mathbb{R} . Is fg uniformly continuous on \mathbb{R} ? If not, give the reason.

(l) A continuous periodic function on \mathbb{R} is bounded and _____ on \mathbb{R} .
(Fill in the blank)

(m) "The derivative of an odd function is an even function." Write true **or** false.

(n) Write the derivative of the function $f(x) = |x|$ for $x \neq 0$.



(o) If f is differentiable on $[a, b]$ and g is a function defined on $[a, b]$ such that $g(x) = kx - f(x)$ for $x \in [a, b]$. If $f'(a) < k < f'(b)$, then find $g'(c)$.

(p) "Suppose $f : [0, 2] \rightarrow \mathbb{R}$ is continuous on $[0, 2]$ and differentiable on $(0, 2)$, with $f(0) = 0$, $f(2) = 1$. If there exists $c \in (0, 2)$, then $f'(c) = \frac{1}{3}$." Is it true or false?

(q) Find $\lim_{x \rightarrow 0} \frac{x^2 + x}{\sin 2x}$.

(r) "The function $f(x) = 8x^3 - 8x^2 + 1$ has two roots in $[0, 1]$." Write true **or** false.

2. Answer **any five** parts : $2 \times 5 = 10$

(a) Use the definition of limit to show that

$$\lim_{x \rightarrow 2} (x^2 + 4x) = 12.$$

(b) Find $\lim_{x \rightarrow 0} x \sin\left(\frac{1}{x^2}\right)$; ($x \neq 0$).

(c) Give an example of a function that has a right-hand limit but not a left-hand limit at a point.

(d) Define $g : \mathbb{R} \rightarrow \mathbb{R}$ by

$$g(x) = \begin{cases} 2x & \text{for } x \in \mathbb{Q} \\ x+3, & \text{for } x \in \mathbb{Q}^c \end{cases}$$

Find all points at which g is continuous.


(e) Show that the 'sine' function is continuous on \mathbb{R} .

(f) Show that the function $f(x) = \frac{1}{x}$ is uniformly continuous on $[a, \infty]$, where $a > 0$.

(g) Using the mean value theorem, show that

$$\frac{x-1}{x} < \ln(x) < x-1 \text{ for } x > 1.$$

(h) Show that $f(x) = x^{1/3}$, $x \in \mathbb{R}$, is not differentiable at $x = 0$.



(i) Let $f(x) = \frac{\ln(\sin x)}{\ln(x)}$

Find $\lim_{x \rightarrow 0^+} f(x)$.

(j) State Darboux's theorem.

3. Answer **any four** parts : $5 \times 4 = 20$

(a) Prove that a number $c \in \mathbb{R}$ is a cluster point of a subset A of \mathbb{R} if and only if there exists a sequence (x_n) in A such that $\lim_{n \rightarrow \infty} x_n = c$ and $x_n \neq c$ for all $n \in \mathbb{N}$.

(b) State and prove squeeze theorem.

(c) Let $A \subseteq \mathbb{R}$, let f and g be functions on A to \mathbb{R} , and let f and g be continuous at a point c in A . Prove that $f-g$ and fg are continuous at c .

(d) Give an example of functions f and g that are both discontinuous at a point c in \mathbb{R} such that $f+g$ and fg are continuous at c .

(e) If $f : A \rightarrow \mathbb{R}$ is a Lipschitz function, then prove that f is uniformly continuous on A .

(f) Determine where the function

$$f(x) = |x| + |x-1|$$

from \mathbb{R} to \mathbb{R} is differentiable and find the derivative.

(g) Find $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x$.

(h) Determine whether or not $x=0$ is a point of relative extremum of the function $f(x) = x^3 + 2$.

4. Answer **any four** parts : $10 \times 4 = 40$

(a) Let $f : A \rightarrow \mathbb{R}$ and let c be a cluster point of A . Prove that the following are equivalent :

(i) $\lim_{x \rightarrow c} f(x) = L$



(ii) Given any ε -neighbourhood $V_\varepsilon(L)$ of L , there exists a δ -neighbourhood $V_\delta(c)$ of c such that if $x \neq c$ is any point $V_\delta(c) \cap A$, then $f(x)$ belongs to $V_\varepsilon(L)$.

(b) (i) Find $\lim_{x \rightarrow 0} \frac{\sqrt{1+2x} - \sqrt{1+3x}}{x+2x^2}$, where $x > 0$. 4

(ii) Prove that $\lim_{x \rightarrow 0} \cos\left(\frac{1}{x}\right)$ does not exist but $\lim_{x \rightarrow 0} x \cos\left(\frac{1}{x}\right) = 0$. 6

(c) (i) Let $f(x) = e^{\frac{1}{x}}$ for $x \neq 0$. Show that $\lim_{x \rightarrow 0^+} f(x)$ does not exist in \mathbb{R} but $\lim_{x \rightarrow 0^-} f(x) = 0$. 5

(ii) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be such that $f(x+y) = f(x) + f(y)$ for all x, y in \mathbb{R} . Suppose that $\lim_{x \rightarrow 0} f(x) = L$ exists. Show that $L = 0$ and then prove that f has a limit at every point c in \mathbb{R} . 5

(d) (i) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by

$$f(x) = \begin{cases} 1 & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

Show that f is not continuous at any point of \mathbb{R} . 5

(ii) Prove that every polynomial function is continuous on \mathbb{R} . 5

(e) Let $A \subseteq \mathbb{R}$, let $f: A \rightarrow \mathbb{R}$, and let $|f|$ be defined by $|f|(x) = |f(x)|$ for $x \in A$. Also let $f(x) \geq 0$ for all $x \in A$ and let \sqrt{f} be defined by $(\sqrt{f})(x) = \sqrt{f(x)}$ for $x \in A$. Prove that if f is continuous at a point c in A , then $|f|$ and \sqrt{f} are continuous at c . 5+5=10

(f) (i) State and prove Bolzano's intermediate value theorem. 1+4=5

(ii) Let A be a closed bounded interval and let $f: A \rightarrow \mathbb{R}$ is continuous on A . Prove that f is uniformly continuous on A . 5



(g) Let $A \subseteq \mathbb{R}$ be an interval, let $c \in A$, and let $f : A \rightarrow \mathbb{R}$ and $g : A \rightarrow \mathbb{R}$ be functions differentiable at c . Prove that

(i) the function $f + g$ is differentiable at c and

$$(f + g)'(c) = f'(c) + g'(c) \quad 5$$

(ii) if $g(c) \neq 0$, then the function $\frac{f}{g}$ is differentiable at c and

$$\left(\frac{f}{g}\right)'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{(g(c))^2} \quad 5$$

(h) State and prove Rolle's theorem. Give the geometrical interpretation of the theorem. (2+5)+3=10

(i) (i) Use Taylor's theorem with $n = 2$ to approximate $\sqrt[3]{1+x}$, $x > -1$. 5

(ii) If $f(x) = e^x$, show that the remainder term in Taylor's theorem converges to zero as $n \rightarrow \infty$ for each fixed x_0 and x . 5

(j) Find the limits :

5+5=10

(i) $\lim_{x \rightarrow 0^+} x^{\sin x}$

(ii) $\lim_{x \rightarrow \frac{\pi}{2}^-} \frac{\tan x}{\sec x}$

